

DISTAL EXTRAHEPATIC BILE DUCT DATA SHEET (ICD10 C24.0)

UICC TNM 9th EDITION STAGING SUMMARY

Stage Group	T stage	N stage	M stage
Stage 0	Tis	N0	M0
Stage I	T1	N0	M0
Stage IIA	T1	N1	M0
	T2	N0	M0
Stage IIB	T2	N1	M0
	T3	N0, N1	M0
Stage IIIA	T1, T2, T3	N2	M0
Stage IIIB	T4	Any N	M0
Stage IV	Any T	Any N	M1

TNM Clinical Classification

T – Primary Tumour

cTX	Primary tumour cannot be assessed
cT0	No evidence of primary tumour
cTis	Carcinoma <i>in situ</i>
cT1	Tumour invades bile duct wall to a depth less than 5 mm
cT2	Tumour invades bile duct wall to a depth of 5 mm up to 12 mm
cT3	Tumour invades bile duct wall to a depth of more than 12 mm
cT4	Tumour involves the coeliac axis, the superior mesenteric artery and/or the common hepatic artery

N – Regional Lymph Nodes

cNX	Regional lymph nodes cannot be assessed
cN0	No regional lymph node metastases
cN1	Metastases to 1–3 regional nodes
cN2	Metastasis to 4 or more regional nodes

M – Distant Metastasis	
cM0	No distant metastasis
cM1	Distant metastasis

TNM Pathological Classification	
<i>The pT and pN categories correspond to the cT and cN categories</i>	
pN0	Histological examination of a regional lymphadenectomy specimen should ordinarily include 12 or more lymph nodes. If the regional lymph nodes are negative, but the number ordinarily examined is not met, classify as pN0