

PERIHILAR BILE DUCTS DATA SHEET (ICD10 C24.0)

UICC TNM 9th EDITION STAGING SUMMARY

Stage Group	T stage	N stage	M stage
Stage 0	Tis	N0	M0
Stage I	T1	N0	M0
Stage II	T2a, T2b	N0	M0
Stage IIIA	T3	N0	M0
Stage IIIB	T4	N0	M0
Stage IIIC	Any T	N1	M0
Stage IVA	Any T	N2	M0
Stage IVB	Any T	Any N	M1

TNM Clinical Classification

T – Primary Tumour

cTX	Primary tumour cannot be assessed
cT0	No evidence of primary tumour
cTis	Carcinoma in situ
cT1	Tumour confined to the bile duct, with extension up to the muscle layer or fibrous tissue
cT2a	Tumour invades beyond the wall of the bile duct to surrounding adipose tissue
cT2b	Tumour invades adjacent hepatic parenchyma
cT3	Tumour invades unilateral branches of the portal vein or hepatic artery
cT4	Tumour invades the main portal vein or its branches bilaterally, or the common hepatic artery, or unilateral second-order biliary radicals with contralateral portal vein or hepatic artery involvement

N – Regional Lymph Nodes

cNX	Regional lymph nodes cannot be assessed
cN0	No regional lymph node metastasis
cN1	Metastases to 1–3 regional lymph nodes
cN2	Metastases to 4 or more regional nodes

M – Distant Metastasis	
-------------------------------	--

cM0	No distant metastasis
------------	-----------------------

cM1	Distant metastasis
------------	--------------------

TNM Pathological Classification	
--	--

<i>The pT and pN categories correspond to the cT and cN categories</i>	
---	--

pN0	Histological examination of a regional lymphadenectomy specimen should ordinarily include 15 more lymph nodes. If the regional lymph nodes are negative, but the number ordinarily examined is not met, classify as pN0
------------	---